

Solar heavy element abundance: constraints from frequency separation ratios of low-degree p modes

William J. Chaplin

School of Physics and Astronomy, University of Birmingham, Edgbaston, Birmingham B15 2TT, U.K.; w.j.chaplin@bham.ac.uk

Aldo M. Serenelli

Institute for Advanced Study, Einstein Drive, Princeton, NJ 08540; aldos@ias.edu

Sarbani Basu

Department of Astronomy, Yale University, P.O. Box 208101, New Haven, CT 06520-8101; sarbani.basu@yale.edu

Yvonne Elsworth

School of Physics and Astronomy, University of Birmingham, Edgbaston, Birmingham B15 2TT, U.K.; y.p.elsworth@bham.ac.uk

Roger New

Faculty of Arts, Computing, Engineering and Sciences, Sheffield Hallam University, Sheffield S1 1WB, U.K.; r.new@shu.ac.uk

and

Graham A. Verner

School of Physics and Astronomy, University of Birmingham, Edgbaston, Birmingham B15 2TT, U.K.; gav@bison.ph.bham.ac.uk

ABSTRACT

We use very precise frequencies of low-degree solar-oscillation modes measured from 4752 days of data collected by the Birmingham Solar-Oscillations Network (BiSON) to derive seismic information on the solar core. We compare these observations to results from a large Monte Carlo simulation of standard solar models, and use the results to constrain the mean molecular weight of the solar core, and the metallicity of the solar convection zone. We find that only

a high value of solar metallicity is consistent with the seismic observations. We can determine the mean molecular weight of the solar core to a very high precision, and, dependent on the sequence of Monte Carlo models used, find that the average mean molecular weight in the inner 20% by radius of the Sun ranges from 0.7209 to 0.7231, with uncertainties of less than 0.5% on each value. Our lowest seismic estimate of solar metallicity is $Z = 0.0187$ and our highest is $Z = 0.0239$, with uncertainties in the range of 12–19%. Our results indicate that the discrepancies between solar models constructed with low metallicity and the helioseismic observations extend to the solar core and thus cannot be attributed to deficiencies in the modeling of the solar convection zone.

Subject headings: Sun: helioseismology - Sun: interior - Sun: abundances

1. Introduction

In a series of papers, Asplund et al. (2004, 2005b, 2005c) and Allende-Prieto et al. (2001, 2002) have revised the spectroscopic determinations of the solar photospheric composition. In particular, their results have determined carbon, nitrogen and oxygen abundances to be lower by about 25% to 35% than previous determinations (Grevesse & Sauval 1998; hereafter GS98). The revision of the oxygen abundance leads to a comparable change in the abundances of neon and argon. Additionally, Asplund (2000) has also determined a somewhat lower value (10%) for the photospheric abundance of silicon compared to the GS98 value. As a result, all the elements for which abundances are obtained from meteoritic measurements have seen their abundances reduced by a similar amount. These measurements have been summarized in Asplund et al. (2005a; hereafter AGS05), and the net result is that the ratio of the mass fraction of heavy elements to hydrogen in the Sun is $Z/X = 0.0165$ (alternatively, $Z = 0.0122$), about 28% lower than the previous value, $Z/X = 0.0229$ ($Z = 0.0169$) given by GS98.

The new low-abundance value for the heavy elements, albeit the result of a much more sophisticated modeling of the solar atmosphere, has given rise to discrepancies between helioseismic observations and predictions from solar models constructed with the low value of Z/X . Solar models constructed with the GS98 composition have shown a remarkable agreement with the solar structure, as determined by helioseismology techniques (Christensen-Dalsgaard et al. 1996; Bahcall et al. 1997; Morel et al. 1999; Basu et al. 2000). However, when the AGS05 composition is adopted in the solar models, the predicted surface helium abundance is too low and the convective envelope too shallow. Additionally, the model sound-speed and density profiles show a degraded agreement with their solar counterparts

when compared to predictions from models that use the older GS98 composition (Bahcall & Pinsonneault 2004; Basu & Antia 2004; Bahcall et al. 2005a, 2005b, 2005c, 2006; Delahaye & Pinsonneault 2006).

The discrepancy between the low- Z/X models and the helioseismic observations has led to attempts to determine the solar metallicity from helioseismic data alone, just as the solar helium abundance was determined by helioseismology. Antia & Basu (2006) used helioseismic data to estimate a value for Z in the solar convection zone of 0.0172 ± 0.002 , i.e., closer to the GS98 value and much larger than the AGS05 value. The uncertainty in their results arose from uncertainties in the equation of state, and a lack of data on acoustic modes of high angular degree ($l > 200$). In this paper we look to the solar core, where uncertainties in the physics of the equation of state and opacities are much lower, to try to constrain the solar metallicity.

We make use of solar p -mode data derived from observations made by the ground-based Birmingham Solar-Oscillations Network (BiSON; Chaplin et al. 1996). The BiSON instruments make disc-averaged observations of the Sun in Doppler velocity. BiSON data can be used to determine very precise frequencies of low- l modes ($l \leq 3$) that can be used to probe the solar core. The possibility of using these data to shed light on the solar abundance problem was explored by Basu et al. (2007). Basu et al. made very specific combinations of the low- l frequencies, the so-called ‘small frequency spacings’ and ‘frequency separation ratios’, to compare models with the observations.

The *small frequency spacings* of the low- l p modes are given by the combination.

$$d_{l+2}(n) = \nu_{n,l} - \nu_{n-1,l+2}, \quad (1)$$

where $\nu_{n,l}$ is the frequency of a mode of degree l and radial order n . The fine spacings are determined predominantly by the sound-speed gradient in the core. Using the asymptotic theory of p modes it can be shown that (see e.g., Christensen-Dalsgaard & Berthomieu 1991)

$$d_{l+2}(n) \simeq -(4l+6) \frac{\Delta_l(n)}{4\pi^2 \nu_{n,l}} \int_0^R \frac{dc}{dr} \frac{dr}{r}, \quad (2)$$

where R is the solar radius, and $\Delta_l(n)$ is the large frequency spacing given by

$$\Delta_l(n) = \nu_{n,l} - \nu_{n-1,l}. \quad (3)$$

The large frequency separation depends inversely on the sound-travel time between the center and the surface of the Sun. The frequencies $\nu_{n,l}$ and $\nu_{n-1,l+2}$ are very similar and hence are affected in a similar way by near-surface effects. By taking this difference in frequency a large part of the effects from the near-surface uncertainties cancels out, making

the spacings a useful probe of the deep solar interior and core. Some residual effects do nevertheless remain. One way of reducing the effects of the near-surface errors is to use the *frequency separation ratios*. The frequency separation ratios (Roxburgh & Vorontsov 2003; Otí Floranes, Christensen-Dalsgaard & Thompson 2005; Roxburgh 2005) are formed from the small frequency spacings and large frequency spacings of the modes. The separation ratios are then constructed according to:

$$r_{02}(n) = \frac{d_{02}(n)}{\Delta_1(n)}, \quad r_{13}(n) = \frac{d_{13}(n)}{\Delta_0(n+1)}. \quad (4)$$

Since both the small and large spacings are affected in a similar manner by near-surface effects, these ratios are somewhat independent of the structure of the surface.

Basu et al. (2007) showed that small spacings and separation ratios for models constructed with the old GS98 composition match the observed BiSON spacings and ratios much more closely than do the spacings and ratios of models with the lower AGS05 composition. In short, models constructed with higher metallicities compare better with the BiSON data than do models constructed with lower metallicities, although the level of agreement deteriorates when the metallicity becomes very large. This indicates that we should be able to determine solar metallicity using the spacing and ratio data.

In this paper, we therefore expand on the work of Basu et al. (2007), and use the small spacings and separation ratios from BiSON data to determine the metallicity of the Sun. We compare the observed spacings and ratios with spacings and ratios of some 12,000 solar models. The models, which were made for an extensive Monte Carlo simulation (Bahcall et al. 2006), account for all the relevant uncertainties entering standard solar model calculations. From this comparison, we show that it is possible to place extremely tight constraints on μ_c , the mean molecular weight averaged over the inner 20 % by radius (i.e., over most of the solar core) of the Sun. Since the mean molecular weight in the core is related to the metallicity at the surface (i.e., the convection zone), we can also place reasonably precise constraints on the heavy element abundance, Z . Both μ_c and Z are measures of solar metallicity, albeit for different regions of the Sun.

The rest of the paper is organized as follows. The observed data and the models used are described in § 2. In § 3, we compare the BiSON separation ratios with the separation ratios of two sequences of models made to test the response of the spacings and ratios to μ_c and Z . In this section we also show how we can obtain a seismic estimate of the solar μ_c and Z by comparison of the BiSON and model spacings and ratios. In § 4 we determine the systematic errors that arise due to uncertainties in the radiative opacities and the equation of state — two quantities are that not amenable to a Monte Carlo type study. In § 5 we expand the BiSON-model comparison by analyzing the results for a grand total of 12,000

Monte Carlo solar models, computed by Bahcall et al. (2006). This analysis allows us to test the impact of changes to several input solar model parameters (in particular changes to the mixture of the heavy elements) on the results. Finally, we summarize our results in § 6.

2. Data and models used

We have made use of Doppler velocity observations made by the BiSON over the 4752-d period beginning 1992 December 31, and ending 2006 January 3. Frequencies were determined by fitting resonant peaks in the power spectrum of the complete time series to yield estimates of the low- l frequencies (e.g., see Chaplin et al. 1999). Prior to calculation of the small spacings and separation ratios, we removed the solar-cycle shifts from the raw fitted low- l frequencies. Details on the process used to remove the solar-cycle shifts can be found in Basu et al. (2007) (see also Chaplin et al. 2005). The BiSON spacings and ratios were then constructed from these corrected frequency data, and the uncertainties on individual fitted frequencies propagated in the usual manner to give the spacing and ratio uncertainties. Table 1 contains those solar-cycle-corrected BiSON mode frequencies which were used to compute the BiSON frequency spacings and separation ratios analyzed in the paper.

Basu et al. (2007) showed that the separation ratios depend on the molecular weight, but they did not determine the exact dependence of the separation ratios on the average mean molecular weight of the core. To do so, we use two very different sets of models. The first set of models, which we refer to as the *test* models, consist of two sequences of ten solar models each. One sequence of models was constructed with the relative heavy element abundances of GS98, while the second sequence was made with the relative heavy abundances of AGS05. To fix the Z/X of a given model in either sequence, the individual relative heavy element abundances of GS98 (or AGS05) were multiplied by the same constant factor. This factor was then changed from one model to another within the sequence. All models in the two sequences were constructed with the same nuclear reaction rates, opacities, equation of state and diffusion rates. These models were constructed to test the dependence of the separation ratios on the *average* mean molecular weight, μ_c and the *total* heavy element abundance, Z . For reference, we include in this paper tables of low- l mode frequencies for two standard models. Table 2 has frequencies for a model with the exact GS98 abundance; while Table 3 has frequencies for a model with the exact AGS05 abundance.

The second set of models comprised a grand total of 12,000 models created for a Monte Carlo study (Bahcall et al. 2006), and we refer to these as the *Monte Carlo* models. The characteristics and methods of computation of the models can be found in Bahcall et al. (2006). Here, we summarize the salient points only.

For each solar model 19 input parameters were drawn randomly from separate probability distributions for every parameter (see Bahcall et al. 2006 for more details). Seven of the input parameters were nuclear reaction rates for important low-energy fusion reactions. The solar age, luminosity, and diffusion coefficient (rate) were the next three parameters. The final nine parameters were the abundances of nine heavy elements: C, N, O, Ne, Mg, Si, S, Ar and Fe. For each solar model, radiative opacity tables corresponding to the randomly generated composition were computed and used. Radiative opacity tables were generated using data and codes provided by the Opacity Project group as presented in Seaton (2005). Low-temperature opacities (for temperatures under 10^4 K) were from Ferguson et al. (2005). Variations of the abundances were dealt with in such a way as to give four sequences of Monte Carlo models. Choices had to be made regarding the underlying mixture, and the probability distribution for the mixture. Two basic mixtures were used: The GS98 and the AGS05 mixtures. Probability distributions were then assigned on the basis of two different estimates of the uncertainties in the abundances of the nine individual elements: ‘conservative’ (large) uncertainties, based on differences between the abundances of the GS98 and AGS05 mixtures; and ‘optimistic’ (small) uncertainties, based on the uncertainties quoted by Asplund et al. (2005a). The content of the four sequences of models may be summarized as follows:

GS-Cons — These 5000 models were made with the ‘conservative’ (large) abundance uncertainties, centered on the GS98 mixture.

GS-Opt — These 1000 models were made with the ‘optimistic’(small) abundance uncertainties, centered on the GS98 mixture.

AGS-Cons — These 1000 models were made with the ‘conservative’ (large) abundance uncertainties, centered on the AGS05 mixture.

AGS-Opt — These 5000 models were made with the ‘optimistic’ (small) abundance uncertainties, centered on the AGS05 mixture.

We have calculated frequencies of low-degree modes for all the models and constructed the small spacings and separation ratios in exactly the same manner as for the observations. The model frequencies come from adiabatic calculations. Since non-adiabatic effects are not included, this leads to a well-known mismatch between the absolute values of the adiabatic model frequencies and the observed frequencies. However, as noted earlier, differences due to these near-surface effects are reduced significantly by taking frequency differences, and using the small spacings and separation ratios.

3. Dependence of the separations on metallicity

In order to parametrize the relation between metallicity and the small spacings and separation ratios, we begin by comparing the BiSON spacings and ratios with the spacings and ratios of the *test* models. To show how the BiSON-model comparisons were made, consider the analysis of the separation ratios. We calculated for each model the differences between the observed BiSON ratios, $r_{02}(n)$ and $r_{13}(n)$, and the model ratios, $r'_{02}(n)$ and $r'_{13}(n)$, i.e.,

$$\Delta r_{l,l+2}(n) = r_{l,l+2}(n) - r'_{l,l+2}(n). \quad (5)$$

These differences were then averaged over n , for each of the $\Delta r_{02}(n)$ and $\Delta r_{13}(n)$, to yield *weighted mean differences*, $\langle \Delta r_{02} \rangle$ and $\langle \Delta r_{13} \rangle$:

$$\langle \Delta r_{l,l+2} \rangle = \frac{\sum_n \Delta r_{l,l+2}(n) / \sigma_{r_{l,l+2}}^2(n)}{\sum_n 1 / \sigma_{r_{l,l+2}}^2(n)}. \quad (6)$$

The formal uncertainties of the BiSON spacings, $\sigma_{r_{l,l+2}}(n)$, were used to weight the averages (with the usual uncertainty-squared Gaussian weighting applied). We averaged data over the ranges where good determinations of the separation ratios were available, here $n = 9$ to 25.

Fig. 1 shows plots of the weighted mean differences of the separation ratios (in %) versus $\ln \mu_c$, the natural logarithm of the average mean molecular weight of the core. The top two panels show data for the GS98 set, and the bottom two panels show data for the AGS05 set. The formal uncertainties on each point, which come from the BiSON data, are not plotted. They are 0.046 % on each $\langle \Delta r_{02} \rangle$; and 0.038 % on each $\langle \Delta r_{13} \rangle$. Fig. 2 shows plots of the weighted mean differences versus the natural logarithm of the surface heavy element abundances, $\ln Z$. Again, the top two panels show data for the GS98 set, and the bottom two panels show data for the AGS05 set.

When a straight line was fitted to the data in each plot (solid lines), the fitting coefficients indicated that a linear dependence was a good model for the data. The fit for the Z data was in all cases described by:

$$\langle \Delta r_{l,l+2} \rangle = \alpha_{l,l+2} + \beta_{l,l+2} \ln Z, \quad (7)$$

where $\alpha_{l,l+2}$ and $\beta_{l,l+2}$ are, respectively, the best-fitting estimates of the intercept and gradient of the straight line. A similar straight-line fitting model was used for the μ_c data, with $\ln \mu_c$ used as the independent variable.

The quality of the Z fits degraded significantly when Z , rather than $\ln Z$, was used as the independent variable. Visible departures from a straight line were then observed. The fits also deteriorated when the fine spacings, rather than the separation ratios, were used. We therefore devote the remainder of the paper to analysis of weighted mean differences made from the separation ratios of the BiSON data and the solar models. These weighted mean differences have then been used to determine the natural logarithm of Z , as opposed to Z itself. We adopted a similar approach to our study of the mean molecular weight in the core. Here, we used the $\ln \mu_c$ of the models as the independent variable for the plots. Again, this was because we found that use of μ_c , rather than $\ln \mu_c$, degraded the quality of the fits somewhat, although not as much as in the case of Z .

With reference to Figures 1 and 2, it is not surprising that the two measures of metallicity affect the separation ratios in similar ways. The quantities μ_c and Z are related, in that a higher Z results in a higher μ_c . Two models with the same Z , nuclear reaction rates, opacities and equation of state can have different values of μ_c only if the diffusion rates are different in the two models. It is, however, not surprising that the dependence of the separation ratios on the two parameters is somewhat different given that μ_c also depends heavily on the helium abundance in the core. All the models are calibrated to have the same radius and luminosity at the solar age, and hence differences in Z generally give rise to differences in the core helium abundance.

3.1. Seismic estimates of solar Z and μ_c

If the observational data are unbiased, the location on the abscissa that marks where each best-fitting straight line passes through zero on the ordinate will give us a ‘seismic’ estimate of the average mean molecular weight μ_c averaged over the inner 20% by radius, and the surface abundance Z of the Sun. (Note that the Z we refer to is always the present-day (solar age) surface Z .) These locations are marked on the various panels of Figures 1 and 2 by the intersecting dotted lines. Conclusions drawn from the test models will of course neglect any dependence of the differences of the separation ratios on changes to other solar model input parameters, including changes to the mixture of the heavy elements. We go on to discuss the impact of such changes, and the overall error budget, in later sections. Here we show explicitly how a value for, and uncertainty on, Z may be estimated from the differences. The same procedures give results for μ_c .

From the best-fitting coefficients, we seek to find $\ln Z$ where $\langle \Delta r_{l,l+2} \rangle = 0$. From Equation 7, we therefore have:

$$\ln Z = -\alpha_{l,l+2}/\beta_{l,l+2},$$

So that

$$Z = \exp[-\alpha_{l,l+2}/\beta_{l,l+2}]. \quad (8)$$

Application of Equation 8 to the $\langle\Delta r_{02}\rangle$ and $\langle\Delta r_{13}\rangle$ data of each set (GS98 or AGS05) will give us four seismic estimates of the solar heavy element abundance. Using GS98 models we get $Z = 0.01798$ and $Z = 0.01774$ for $\langle\Delta r_{02}\rangle$ and $\langle\Delta r_{13}\rangle$ respectively. The corresponding results using the AGS05 models are $Z = 0.01617$ and $Z = 0.01611$. A similar analysis of μ_c results in values of 0.7253 and 0.7244 when GS98 models are used, and 0.7260 and 0.7255 when AGS05 test models are used. These estimates are listed in the fourth column of Tables 4 and 5, along with estimates of the goodness-of-fit and uncertainties.

We make use of the observed scatter (variance) of the differences $\langle\Delta r_{02}\rangle$ and $\langle\Delta r_{13}\rangle$ about their best-fitting straight line to estimate the uncertainty on the seismic estimates of solar μ_c and Z . In short, we translate the characteristic scatter on the ordinate into an implied uncertainty on the abscissa. To determine the uncertainty in Z (the same procedure was used for μ_c) we first determine the set of residuals about the best-fitting straight line, i.e., for each point we compute

$$\delta\langle\Delta r_{l,l+2}\rangle = \langle\Delta r_{l,l+2}\rangle - [\alpha_{l,l+2} + \beta_{l,l+2} \ln Z].$$

The variance of these residuals yields an estimate of their 1σ standard deviation, which we call $\sigma_{\langle\Delta r_{l,l+2}\rangle}$. This characteristic uncertainty on the residuals may be translated into an implied uncertainty on $\ln Z$ via the best-fitting gradient, i.e.,

$$\sigma(\ln Z) = \sigma_{\langle\Delta r_{l,l+2}\rangle}/\beta_{l,l+2}. \quad (9)$$

The equivalent 1σ limits on Z are then bounded by $\exp[\ln Z + \sigma(\ln Z)]$ and $\exp[\ln Z - \sigma(\ln Z)]$. Our estimate of the uncertainty is itself uncertain through the uncertainty on the gradient, $\beta_{l,l+2}$. Thus we require that N be large enough to ensure that the best-fitting gradient (and our ‘look-up curve’) is well constrained. Here, gradients for the Z fits were returned to a fractional precision of better than 1%.

The third column of Tables 4 and 5 shows the computed $\sigma_{\langle\Delta r_{l,l+2}\rangle}$ (in %). The fifth column gives the implied 1σ uncertainties on μ_c and Z , which we call $\sigma(\mu_c)$ and $\sigma(Z)$. Here, the positive and negative uncertainties were the same, at the level of precision of the data, because the fractional uncertainties were so small. The sixth column shows the implied precision in the determination of μ_c and Z (in %). For each of the GS98 and AGS05 sets, we also combined the estimates from $\langle\Delta r_{02}\rangle$ and $\langle\Delta r_{13}\rangle$ – on the assumption the estimates are independent – to give the estimates shown in the third and sixth rows of the tables.

Inspection of the μ_c results in Table 4 shows that all results (both individual and combined) are consistent with one another. This is not surprising given the almost direct dependence of the sound speed, and its derivative, on μ_c . The combined estimate obtained with

the GS98 models differs from that obtained with the AGS05 models by only 1σ . As we shall see in § 5, the uncertainties on the seismic estimates of μ_c do not increase much when other changes to the solar model input parameters are considered.

Inspection of the Z results in Table 5 shows that the individual, and combined, estimates for solar Z are all significantly higher than the ‘low’ Asplund et al. value of $Z \sim 0.0122$. However, our combined GS estimate (0.01785) and its combined AGS counterpart (0.01611) differ from each other by $\approx 16\sigma$ (combined uncertainty). This difference might at first glance be seen as a cause for concern and an indication that systematic errors are much larger than the random errors caused by uncertainties in the observed frequencies. However, we go on to show in § 5 that other systematic sources of error – arising from the sensitivity of the separation ratios to other parameters of the solar models, including the relative mixture of the heavy elements – mean that realistic estimates of the uncertainties on Z are actually larger in size than the uncertainties given in Table 5. This is in stark contrast to what is found for μ_c (see previous paragraph).

Disagreement between the seismic results obtained from the GS98 and AGS98 models can to a large extent be understood in terms of the differences in the two mixtures. The influence of the relative mixture is clear from the fact that the separation ratios for the AGS05 models and the GS98 models are different for the same value of Z ; the differences are less pronounced for μ_c . This finding is not difficult to understand. For the calibrated solar models used in this work, the dominant contribution to the Z and μ_c of each model comes from different elements. For Z , the dominant elements, in order of importance, are oxygen, carbon, neon and nitrogen. The value of μ_c is determined by the mass fractions of helium and hydrogen in the core. The abundances of hydrogen and helium in the core depend strongly on the abundances of heavy metals that contribute to the opacity in the core. These metals, again in order of importance, are iron, silicon, sulfur and oxygen. The difference between the GS98 and AGS05 mixture lies predominantly in the relative abundances of oxygen, carbon, nitrogen and neon, and much less so in the abundances of iron, silicon and sulfur. This explains why for the same Z , μ_c is different for the GS98 and AGS05 models, as can be seen from Fig. 3.

Since the separation ratios depend basically on μ_c and temperature, the location at which $\langle \Delta r_{l,l+2} \rangle = 0$ occurs at slightly different values for the two sets of models. The slopes of the $\langle \Delta r_{l,l+2} \rangle$ - $\ln Z$ curves for the GS98 and AGS05 models (Figures 2) are also different for the same reason. We investigate other sources of systematic errors in § 5. The results in § 5 show that realistic estimates of the uncertainties on Z are actually larger in size than the uncertainties given in Table 5.

4. Uncertainties due to opacity and equation of state

The results obtained with the test models have two obvious limitations. First, they do not test the impact of changes to the relative mixture of the heavy elements, except to reveal that changes in relative abundances matter. And second, the models test the dependence of the separation ratios on μ_c and Z for a *fixed* set of solar model input parameters, with only the Z/X varied. The study above does not deal with uncertainties in the seismic estimates of solar μ_c and Z caused by other input parameters, such as nuclear reaction rates, diffusion rates, uncertainties in the relative mixtures, etc. We investigate these effects by conducting a Monte Carlo study, which is described below in § 5. This study allowed us to test the impact of 19 solar model input parameters on the results. There are, however, two important inputs that are not amenable to a Monte Carlo study, and these are radiative opacities and the equation of state (EOS). These two quantities cannot be described by a single number and hence we are forced to use a different approach to determine the uncertainties in the separation ratios caused by uncertainties in opacities and EOS. We use an approach similar to that used by Bahcall et al. (2006).

We determined the uncertainty introduced in $\langle \Delta r_{l,l+2} \rangle$ by the opacities as follows. We computed a pair of solar models with the same input parameters and EOS, but one was made with opacities from the OP project (Badnell et al. 2005) and the other with opacities from OPAL (Iglesias & Rogers 1996). For this matched pair of models we get $\langle \Delta r_{l,l+2} \rangle_{\text{OP},i}$ and $\langle \Delta r_{l,l+2} \rangle_{\text{OPAL},i}$ (here the subscript i denotes the pair of matched models). The unbiased estimator for the variance of the difference is

$$s_i^2(\langle \Delta r_{l,l+2} \rangle (\text{opacity})) = \left[\langle \Delta r_{l,l+2} \rangle_{\text{OP},i} - \langle \Delta r_{l,l+2} \rangle_{\text{OPAL},i} \right]^2 / 2. \quad (10)$$

and we adopt this quantity as the standard deviation squared, $\sigma_i^2(\langle \Delta r_{l,l+2} \rangle (\text{opacity}))$. In order to obtain a more representative value for $\sigma_{\text{opac}}(\langle \Delta r_{l,l+2} \rangle)$ we averaged this difference over a set of $N = 40$ matched pairs of models, where the 19 other input parameters were varied for different matched pairs. The final expression we adopt for $\sigma_{\text{opac}}(\langle \Delta r_{l,l+2} \rangle)$ is

$$\sigma_{\text{opac}}(\langle \Delta r_{l,l+2} \rangle) = \sqrt{N^{-1} \sum_i s_i^2(\langle \Delta r_{l,l+2} \rangle (\text{opacity}))}. \quad (11)$$

An analogous procedure was used for the EOS, but in this case one model in each pair was computed using the 2001 OPAL EOS (Rogers 2001, Rogers & Nayfonov 2002), while the other model was computed using the 1996 OPAL EOS (Rogers et al. 1996).

In addition to the uncertainties on $\langle \Delta r_{02} \rangle$ and $\langle \Delta r_{13} \rangle$ given by Equation 11, we applied the same procedure to compute the implied uncertainties for the values of μ_c and Z predicted

by the solar models. Uncertainties in the opacity and EOS will affect the solar model results for Z and μ_c , i.e. Z and μ_c will be somewhat different for the two models in each of the matched pairs described above. The results are given in Table 6.

The impact on the separation ratios of uncertainty in the radiative opacities is easy to understand — changes in opacity cause changes in temperature, which in turn change sound speed and its derivative, thereby changing the separation ratios. However, since the opacity uncertainties are small in the core, the overall effect is quite modest. The impact of uncertainty in the EOS is more important. This might seem surprising, until one realizes that the 1996 OPAL EOS did not treat relativistic effects properly at temperatures and densities relevant to the solar core. This results in a somewhat deficient core structure (see Elliott & Kosovichev, 1998). Since the deficiency is mainly in the core, it will affect the low- l separations used in this work disproportionately and cause larger uncertainties in the seismic estimates of solar Z and μ_c . The updated 2001 OPAL EOS has the correction put in. Thus the uncertainty in the seismic estimates of solar Z and μ_c caused by EOS uncertainties may be considered to be an upper limit to the EOS effects.

5. The Monte Carlo study

In this section we present results, using extended sequences of solar models, which seek to address the influence of other input parameters on the seismic solar Z results reported in § 3.1. The bulk of the results come from tests on 12,000 solar models created for a Monte Carlo study of the dependence of solar model characteristics on different input parameters (see Bahcall et al. 2006). As discussed in § 2, the models have 19 different input parameters selected at random from a distribution of the inputs.

Scatter plots of the weighted mean differences of the separation ratios (in %) versus the natural logarithm of μ_c are shown in Figures 4 and 5 for all four sequences of Monte Carlo models. Figures 6 and 7 show the corresponding plots against $\ln Z$. Uncertainties caused by uncertainties in the radiative opacities and EOS have been included by adding to the data random components with normal distributions characterized by the standard deviations given in Table 6. The solid lines in each panel are the best-fitting straight lines for the data. The dotted lines intersect at the location along each best-fitting line where the weighted mean difference is zero.

We again adopted a linear model for the analysis in the light of: (i) the results on the test models (discussed in Section 3 above); and (ii) the observed scatter on the plots, which precluded us from imposing a more complicated fitting model. Detailed breakdowns of fitting

results for the Monte Carlo sequences are presented in Table 7 (for μ_c), and Table 8 (for Z).

The main results obtained from the Monte Carlo sequences are as follows:

1. Before turning to discussion of the results on μ_c and Z , we consider first the $\langle\Delta r_{02}\rangle$ and $\langle\Delta r_{13}\rangle$ of the model sequences. In the case of the AGS-Opt sequence, the distribution of $\langle\Delta r_{02}\rangle$ is characterized by a mean value of -2.05% and a standard deviation of 0.75% . For $\langle\Delta r_{13}\rangle$ the corresponding values are, respectively, -1.80% and 0.54% . These values imply a difference with helioseismology measurements of 2.7σ and 3.3σ for $\langle\Delta r_{02}\rangle$ and $\langle\Delta r_{13}\rangle$ respectively. Note from Table 6 that the EOS has a large impact on the total standard deviations of the $\langle\Delta r_{02}\rangle$ and $\langle\Delta r_{13}\rangle$ distributions. As discussed previously, we are probably overestimating the EOS uncertainties in the separation ratios. Consequently, the 2.7 and $3.3\text{-}\sigma$ differences should be considered as robust upper limits to the real discrepancy.

On the other hand, for the GS-Opt sequence the mean and standard deviation of the $\langle\Delta r_{02}\rangle$ distribution are -0.43% and 0.76% respectively, while for $\langle\Delta r_{13}\rangle$ we get -0.32% and 0.53% for the mean and the standard deviation. These numbers translate into differences of only $\approx 0.6\sigma$ with the helioseismology measurements. It is worth mentioning that we perform this comparison only for Monte Carlo sequences having the optimistic choice of uncertainties, because the aim is to compare helioseismology measurements with solar models that adopt compositions (central values and uncertainties) given by the solar abundance determinations, i.e. GS98 and AGS05.

2. As in the case of results obtained using the test models, the average mean molecular weight in the inner 20% by radius, μ_c is determined to much higher precision than Z (results in Table 7). The precision in each of the combined measures of μ_c is better than 0.5% . Even with the improved precision, the estimates given by analysis of the four sequences of models are in excellent agreement with one another and provide a very robust determination of μ_c . The largest difference between any of the two combined measures is significant at only $\sim 1\sigma$. The average of our four combined seismic estimates is $\mu_c = 0.7226$.
3. From the AGS-Opt sequence of models, i.e. that adopting the AGS05 central values and uncertainties for the composition, we find that solar models constructed with the AGS05 composition have an average μ_c of 0.7088 ± 0.0029 (1σ uncertainty). This value differs from our combined seismic estimates of solar μ_c , which are given in Table 7, by between 3.1 (for the combined AGS-Cons measure) and 3.7σ (for the combined AGS-Opt measure) (where σ is determined by adding in quadrature the uncertainties from the observations and the uncertainty in the theoretical distribution). On the other

hand, in the case of models adopting the GS98 composition, we derive an average value of $\mu_c = 0.7203 \pm 0.0029$ for the optimistic (i.e., lower) uncertainties on the abundance inputs. All the combined seismic estimates agree with this value to well within 1σ .

4. All seismic estimates of solar Z are high (see Table 8). They lie noticeably above the low Asplund et al. (2005a) (AGS05) value. This is clear even from visual inspection of the plots in Fig. 7, which shows results for models with heavy-element mixtures based on the AGS05 values. The location on the abscissa where the best-fitting lines pass through zero lie well to the high- Z side of each cloud of points. None of the AGS-Opt models has a mean weighted difference greater than zero. Estimation of Z from these data therefore amounts to an *extrapolation*, rather than the *interpolation* as is possible with the GS-Cons and GS-Opt models.

From the AGS-Opt sequence of models, which has abundance uncertainties consistent with Asplund et al. (2005a), we find that solar models constructed with the AGS05 composition have an average Z of 0.0125 ± 0.0007 (1σ uncertainty). This value differs from our combined seismic estimates of solar Z , which are given in Table 8, by between 2.0σ (for the combined GS-Cons measure) and 4.3σ (for the combined AGS-Opt measure) (where σ is determined by adding in quadrature the uncertainties from the observations and the uncertainty in the theoretical distribution).

5. All seismic estimates of solar Z are slightly higher than the Grevesse & Sauval (1998) (GS98) value, of $Z = 0.0169$ (see Table 8). This is clear from visual inspection of the plots in Fig. 6, which shows results for models with heavy-element mixtures based on the GS values. The best-fitting lines pass through zero in the high- Z parts of the distributions of points. Differences between the GS98 value and our combined seismic estimates of solar Z lie between 0.5σ (for the combined GS-Cons measure) and 2.0σ (for the combined AGS-Opt measure).
6. All seismic estimates of solar Z in Table 8 are in good agreement with the seismic estimate of Antia & Basu (2006), which was $Z = 0.0172 \pm 0.0020$. The Antia & Basu result was quoted with a precision of just over 12%. The precision in our four, combined measures of Z (rows 3, 6, 9 and 12 of Table 8) ranges from $\sim 12\%$ to $\sim 19\%$. The largest difference between any of the two combined measures is significant at only $\sim 1\sigma$.

The observed scatter in Figures 4 to 7 deserves discussion, as do the differences between the results given by the Monte Carlo sequence of models and the test models from § 3.

The scatter in the computed $\langle \Delta r_{02} \rangle$ and $\langle \Delta r_{13} \rangle$ about the best-fitting straight lines – as characterized by the $\sigma_{\langle \Delta r_{l,l+2} \rangle}$ in column 3 of Tables 7 and 8 – far exceeds the sizes

of the formal uncertainties on the mean differences found in § 3. Recall that the formal uncertainty (a result of uncertainties in the BiSON data) is only 0.046 % on each value of $\langle \Delta r_{02} \rangle$; and 0.038 % on each value of $\langle \Delta r_{13} \rangle$. The uncertainties on the seismic estimates of solar Z and μ_c computed from analysis of the Monte Carlo data are therefore dominated by scatter introduced by the input parameter choices for the solar models. As we discuss below, the largest contribution to this scatter comes from the relative abundance of heavy elements that contribute to Z . It is worth pointing out that while the uncertainties on the seismically estimated values of solar Z , obtained with the Monte Carlo sequences, are much larger (by over an order of magnitude) than the uncertainties obtained from the test models, corresponding differences in the uncertainties on the seismic values of μ_c , i.e., $\sigma(\mu_c)$, are just a few times larger.

The 1σ standard deviation of the fitting residuals, $\sigma_{\langle \Delta r_{l,l+2} \rangle}$, changes depending on whether ‘conservative’ (large) or ‘optimistic’ (small) uncertainties are used for the input abundances to the solar models. The $\sigma_{\langle \Delta r_{l,l+2} \rangle}$ values for the GS-Cons and AGS-Cons sequences are between ~ 20 and ~ 60 % higher than the corresponding values of the GS-Opt and AGS-Opt sequences. This is not surprising — the ‘conservative’ GS-Cons and AGS-Cons sequences cover, by their very nature, a large, ‘pessimistic’ range of input abundance values, thus the scatter due to the uncertainties in the relative abundance is also larger than that for the ‘optimistic’ sets. The uncertainties that arise from using the GS-Cons and the AGS-Cons models may therefore be treated as respectable *upper* bounds for $\sigma(Z)$ and $\sigma(\mu_c)$.

The gradients and intercepts of the best-fitting lines of the relations between the separation ratios and the metallicity of the test models (Figure 2) are steeper and higher, respectively, than those of their Monte-Carlo counterparts (Figures 6 and 7). The gradients are in some cases steeper by as much as a factor of 3. There are also differences between one Monte Carlo sequence and another, with the gradients differing by up to ~ 35 %; the gradients are constrained typically to much higher precision. However, in spite of these differences, the zero-crossing points, which serve to provide the estimates of solar Z , return consistently robust values.

Again, it is not very difficult to explain why the gradients are higher for the sets of test models compared to the sets of Monte Carlo models, and why the gradients differ from one Monte Carlo set to another. The answer again lies in the impact of changes to the relative mixture of heavy elements. Each set of linear test models has the same relative mixture of elements; these models represent the case of perfect correlation between all the abundance uncertainties. A given value of Z gives a unique value of μ_c , and other conditions in the core (e.g., temperature), and thus matches to a unique value of $\langle \Delta r_{02} \rangle$ and $\langle \Delta r_{13} \rangle$. The Monte Carlo sequences simulate the opposite situation; here, all the abundance

uncertainties are assumed to be independent of one another. The different dependences on relative abundances of Z and μ_c imply that their variations in the MC sequences become, to some degree, uncorrelated. For example, large changes in CNO elements give rise to large changes in Z but have a much more modest impact on μ_c (and consequently on the separation ratios). The result is that in the MC sequences the one-to-one relation between Z , μ_c and other conditions in the core is lost, as is the one-to-one correlation between Z and $\langle\Delta r_{02}\rangle$ or $\langle\Delta r_{13}\rangle$. The overall effect is to force the linear fitting gradients to shallower values when a range of mixtures is admitted in the models. There are changes in the details of the mixtures between the different Monte Carlo sequences, and hence some (albeit much more modest) changes in gradient are seen from one sequence to another.

It is worth adding that changes to the diffusion rates do not alter significantly the fitting gradients. If one takes appropriate subsets of the Monte Carlo sequences, one finds that the relation between the rate of diffusion and μ_c is always the same. And the diffusion rate and Z appear to be only very weakly correlated. Changes to the diffusion do not affect the relation between the fitting gradients of $\langle\Delta r_{02}\rangle$ or $\langle\Delta r_{13}\rangle$ versus $\ln Z$ or $\ln \mu_c$. The only effects seen are changes to the intercept of the fits between the separation ratios and the metallicities, which change the seismic estimates of Z , and thus diffusion is a relevant source of uncertainty. We find that physical inputs other than the relative mixture, the diffusion coefficients, and the EOS have a much smaller effect on the uncertainties of the estimated values of solar μ_c and Z .

6. Results and Discussion

We have used the frequencies of low-degree acoustic oscillations of the Sun, determined by the BiSON network over a period of 4752 days, to try and determine the metallicity Z of the Sun. We did so by comparing the frequency separation ratios of a large set of solar models, from four different Monte Carlo sequences, with the BiSON observations. Specifically, we used a weighted average of the difference of the separation ratios. We find that in addition to giving good constraints on the solar metallicity, which by definition is the abundance of heavy elements in the solar convection zone, the comparison provides an excellent means to determine the mean molecular weight of the solar core.

The frequency separation ratios of the low-degree acoustic modes are sensitive to the conditions in the solar core. By using the weighted mean differences of the separation ratios, we have obtained seismic estimates for the mean molecular weight of the solar core (μ_c). The seismic estimates are robust and depend only very weakly on the solar models used to construct the weighted mean differences. All of the four sequences of Monte Carlo models

used in this work allow us to determine μ_c to a precision of better than 0.5%, and estimates from different sequences are consistent with each other to better than 1σ . This is true even for the seismic estimates obtained using solar models with the AGS05 composition. These solar models do not agree with helioseismic results on the solar sound-speed and density profiles, surface helium abundance, depth of the convective envelope and frequency separation ratios. We have now shown that solar models constructed with the AGS05 composition (central values and uncertainties) fail to reproduce the seismically determined average mean molecular weight of the solar core by more than 3σ . On the other hand, solar models with the older GS98 composition have μ_c well within 1σ of our seismic estimate of μ_c in the Sun.

We get estimates for the solar metallicity in the range between $Z = 0.0187$ and $Z = 0.0229$ for the four Monte Carlo sequences with uncertainties in the range of 12% to 19% on each measurement. All our seismic estimates for metallicity are consistent with each other at about the 1σ level, and are higher than the solar metallicity derived by Asplund et al. (2005) by between 2.1 and 4.3σ . Our estimates are also consistent with the seismically derived value of $Z = 0.0172 \pm 0.0020$ obtained by Antia & Basu (2006). Finally, our estimates are slightly higher (by between 0.5 and 2σ) than the solar metallicity value recommended by Grevesse & Sauval (1998).

The Antia & Basu (2006) results were obtained by looking at the near-surface ionization zones. The signature of interest was the change in the adiabatic index resulting from ionization of material, which depends on the equation of state and the metallicity. The biggest source of systematic error in the results of Antia & Basu was therefore the equation of state. Here, we have obtained very similar results for solar Z by looking at a region where very different physical inputs matter. The main source of systematic error for this study was the relative abundance of the heavy elements. Thus errors in the physical inputs used to construct the solar models are not the reason why we obtain consistency high estimates of solar Z . As a matter of fact, even when we use low- Z models, we still get estimates of solar Z that are high.

When the AGS05 abundances were published, it was soon noticed that solar models constructed with those abundances did not agree well with the helioseismically determined properties of the Sun. In particular, the most obvious discrepancy was in the position of the convection-zone base. It was speculated that this implied that the physical inputs to the models, particularly the opacities, were incorrect. The fact that the helium abundance in the convection zone was also incorrect lead to the belief that the problems are localized to the outer parts of the Sun. Based on the results of this paper, we conclude that this is not the case and that the problems extend to the core. It therefore seems unlikely that the origin of the discrepancy lies in the simplified modeling of the regions close to the tachocline

or in the treatment of convection which is adopted in standard solar model calculations. While rotation and its associated mixing are certainly not modeled in the standard solar models used in this work, it should be noted that solar models that attempt to account for rotation (and some degree of rotationally induced mixing) predict a lower helium abundance in the solar core, which translates into a lower average mean molecular weight (Palacios et al. 2006). Our results seem to indicate that this would make the disagreement between models with the AGS05 composition and helioseismology even worse.

This paper utilizes data collected by the Birmingham Solar-Oscillations Network (BiSON), which is funded by the UK Science Technology and Facilities Council (STFC). We thank the members of the BiSON team, colleagues at our host institutes, and all others, past and present, who have been associated with BiSON. GAV acknowledges the support of STFC. SB acknowledges partial support from NSF grant ATM-0348837. AMS is partially supported through a John Bahcall Membership and a Ralph E. and Doris M. Hansmann Membership at the IAS.

REFERENCES

- Allende Prieto, C., Lambert, D. L., & Asplund, M. 2001, *ApJ*, 556, L63
- Allende Prieto, C., Lambert, D. L., & Asplund, M. 2002, *ApJ*, 573, L137
- Antia, H.M., Basu, S. 2006, *ApJ*, 644, 1292
- Asplund, M. 2000, *A&A*, 359, 755
- Asplund, M., Grevesse, N., Sauval, A. J., Allende Prieto, C., & Kiselman, D. 2004, *A&A*, 417, 751
- Asplund, M., Grevesse, N., & Sauval, A. J. 2005a, in *ASP Conf. Ser. 336, Cosmic Abundances as Records of Stellar Evolution and Nucleosynthesis*, ed. T. G. Barnes III, & F. N. Bash (San Francisco: ASP), 25
- Asplund, M., Grevesse, N., Sauval, A. J., Allende Prieto, C., Blomme, R., 2005b, *A&A*, 431, 693
- Asplund, M., Grevesse, N., Sauval, A. J., Allende Prieto, C., & Kiselman, D. 2005c, *A&A*, 435, 339

- Badnell, N. R., Bautista, M. A., Butler, K., Delahaye, F., Mendoza, C., Palmeri, P., Zeippen, C. J., & Seaton, M. J. 2005, MNRAS, 360, 458
- Bahcall, J. N., Basu, S., Pinsonneault, M. H., & Serenelli, A. M. 2005a, ApJ, 618, 1049
- Bahcall, J. N., Basu, S., Serenelli, A.M. 2005b, 631, 1281
- Bahcall, J.N., Pinsonneault, M.H. 2004, Phys. Rev. Lett., 92, 121301
- Bahcall, J.N., Pinsonneault, M.H., Basu, S., Christensen-Dalsgaard, J. 1997, Phys. Rev. Lett., 78, 171
- Bahcall, J. N., Serenelli, A. M., & Basu, S. 2005c, ApJ, 621, L85
- Bahcall, J.N., Serenelli, A.M., Basu, S., 2006, ApJSS, 165, 400
- Basu, S., & Antia, H. M. 2004, ApJ, 606, L85
- Basu, S., Chaplin, W. J., Elsworth, Y. New, R., Serenelli, A. M., Verner, G. A., 2007, ApJ, 655, 660
- Basu, S., Pinsonneault, M.H., Bahcall, J.N. 2000, ApJ, 529, 1084
- Chaplin W. J., Elsworth Y., Howe R., Isaak G. R., McLeod C. P., Miller B. A., New R., van der Raay H. B. & Wheeler S. J., 1996, Sol. Phys., 168, 1
- Chaplin W. J., Elsworth Y., Isaak G. R., Miller B. A., New R., 1999, MNRAS, 308, 424
- Chaplin W. J., Elsworth Y., Miller B. A., New R., Verner G. A., 2005, ApJ, 635, L105
- Christensen-Dalsgaard, J., Berthomieu, G. 1991, in Solar Interior and Atmosphere, eds. A.N. Cox, W.C.Livingston, M.S. Matthews, University of Arizona Press, Tucson, p401
- Christensen-Dalsgaard, J., Däppen, W., Ajukov, S.V., et al. 1996, Science, 272, 1286
- Delahaye, F., & Pinsonneault M. H. 2006, ApJ, 649, 529
- Elliott, J.R., Kosovichev, A.G. 1998, ApJ, 500, L199
- Ferguson, J. W., Alexander, D. R., Allard, F., Barman, T., Bodnarik, J. G., Hauschildt, P. H., Heffner-Wong, A., & Tamanai, A. 2005, ApJ, 623, 585
- Grevesse, N., & Sauval, A. J. 1998, in Solar composition and its evolution — from core to corona, eds., C. Fröhlich, M. C. E. Huber, S. K. Solanki, & R. von Steiger, Kluwer, Dordrecht, p. 161

- Iglesias, C. A., & Rogers, F. J. 1996, ApJ, 464, 943
- Morel, P., Pinchon, B., Provost, J., Berthomieu, G. 1999, A&A, 350, 275
- Otí Floranes H., Christensen-Dalsgaard J., Thompson M. J., 2005, MNRAS, 356, 671
- Palacios, A., Talon, S., Turck-Chieze, S., & Charbonnel, C. 2006, to appear in Proceedings of the SOHO 18 / GONG 2006 / Workshop (ESA SP-624). “Beyond the Spherical Sun: a new era for helio- and asteroseismology”. 7-11 August, 2006. Sheffield, UK. Editor: K.Fletcher. astro-ph/0609381.
- Rogers, F. J. 2001, Contrib. Plasma Phys., 41, 179
- Rogers, F. J., Swenson, F. J., & Iglesias, C. A. 1996, ApJ, 456, 902
- Rogers, F. J., & Nayfonov, A. 2002, ApJ, 576, 1064
- Roxburgh I. W., Vorontsov S. V., 2003, A&A, 411, 215
- Roxburgh I. W., 2005, A&A, 434, 665
- Seaton, M. J., 2005, MNRAS, 362, L1
- Seaton, M. J., Badnell, N. R., 2004, MNRAS, 354, 457

Table 1. Solar-cycle corrected BiSON frequencies (in μHz) used in the paper

n	$l = 0$	$l = 1$	$l = 2$	$l = 3$
8		1329.629 ± 0.004	1394.682 ± 0.011	1450.986 ± 0.038
9	1407.481 ± 0.006	1472.841 ± 0.005	1535.861 ± 0.008	1591.575 ± 0.014
10	1548.343 ± 0.008	1612.717 ± 0.006	1674.527 ± 0.008	1729.109 ± 0.017
11	1686.588 ± 0.012	1749.296 ± 0.008	1810.323 ± 0.012	1865.281 ± 0.019
12	1822.212 ± 0.014	1885.089 ± 0.010	1945.804 ± 0.017	2001.218 ± 0.022
13	1957.432 ± 0.015	2020.798 ± 0.014	2082.105 ± 0.018	2137.781 ± 0.025
14	2093.496 ± 0.016	2156.781 ± 0.017	2217.678 ± 0.022	2273.521 ± 0.031
15	2228.768 ± 0.018	2291.980 ± 0.014	2352.196 ± 0.031	2407.660 ± 0.034
16	2362.767 ± 0.025	2425.587 ± 0.019	2485.856 ± 0.026	2541.677 ± 0.032
17	2496.140 ± 0.022	2559.196 ± 0.022	2619.670 ± 0.025	2676.191 ± 0.031
18	2629.656 ± 0.021	2693.332 ± 0.021	2754.454 ± 0.024	2811.352 ± 0.029
19	2764.128 ± 0.021	2828.097 ± 0.021	2889.545 ± 0.024	2946.981 ± 0.029
20	2899.010 ± 0.019	2963.306 ± 0.020	3024.687 ± 0.024	3082.319 ± 0.035
21	3033.736 ± 0.021	3098.127 ± 0.022	3159.800 ± 0.028	3217.712 ± 0.040
22	3168.612 ± 0.025	3233.147 ± 0.026	3295.087 ± 0.035	3353.387 ± 0.054
23	3303.537 ± 0.030	3368.495 ± 0.031	3430.807 ± 0.047	3489.430 ± 0.071
24	3439.006 ± 0.042	3504.195 ± 0.040	3567.005 ± 0.057	3626.022 ± 0.101
25	3574.896 ± 0.055	3640.347 ± 0.052		
26	3710.942 ± 0.089			

Table 2. Model frequencies (in μHz) for exact GS98 abundance

n	$l = 0$	$l = 1$	$l = 2$	$l = 3$
8		1329.356	1394.304	1450.684
9	1407.157	1472.596	1535.579	1591.042
10	1548.061	1612.210	1674.187	1728.762
11	1686.350	1748.958	1809.846	1864.909
12	1821.823	1884.864	1945.720	2001.096
13	1957.418	2020.779	2082.327	2138.317
14	2093.866	2157.501	2218.607	2274.732
15	2229.760	2293.493	2354.395	2410.169
16	2365.099	2428.289	2489.179	2545.531
17	2499.604	2563.249	2624.244	2681.130
18	2634.334	2698.557	2760.298	2817.711
19	2770.118	2834.665	2896.607	2954.613
20	2906.128	2971.186	3033.263	3091.349
21	3042.414	3107.438	3169.828	3228.318
22	3178.697	3243.930	3306.347	3365.269
23	3314.908	3380.526	3443.372	3502.561
24	3451.650	3517.369	3580.551	3640.271
25	3588.594	3654.679		
26	3725.807			

Table 3. Model frequencies (in μHz) for exact AGS05 abundance

n	$l = 0$	$l = 1$	$l = 2$	$l = 3$
8		1328.009	1392.406	1448.604
9	1405.414	1470.904	1533.686	1588.631
10	1546.296	1609.957	1671.676	1726.210
11	1684.076	1746.706	1807.136	1861.847
12	1819.239	1882.140	1943.031	1998.140
13	1954.946	2018.007	2079.174	2135.089
14	2090.926	2154.618	2215.457	2271.133
15	2226.755	2290.126	2350.930	2406.628
16	2361.873	2425.004	2485.506	2541.695
17	2496.103	2559.770	2620.763	2677.364
18	2631.043	2695.017	2756.563	2813.952
19	2766.599	2831.216	2892.902	2950.568
20	2902.576	2967.404	3029.432	3087.386
21	3038.792	3103.708	3165.737	3224.098
22	3174.772	3240.032	3302.407	3360.995
23	3311.132	3376.466	3439.208	3498.371
24	3447.692	3513.433	3576.329	3635.823
25	3584.526	3650.497		
26	3721.747			

Table 4. μ_c results for ‘linear test’ models

Set	N	$\sigma_{\langle \Delta r_{l,l+2} \rangle}$ (%)	μ_c	$\sigma(\mu_c)$	$\sigma(\mu_c)/\mu_c$ (%)	χ^2_{N-2}
GS-Lin r_{02}	10	0.136	0.7253	± 0.0012	0.17	9.9
GS-Lin r_{13}	10	0.099	0.7244	± 0.0010	0.14	7.4
GS-Lin comb.			0.7248	± 0.0008	0.11	
AGS-Lin r_{02}	10	0.108	0.7260	± 0.0009	0.12	6.1
AGS-Lin r_{13}	10	0.089	0.7255	± 0.0008	0.11	6.1
AGS-Lin comb.			0.7258	± 0.0006	0.08	

Table 5. Z results for ‘linear test’ models

Set	N	$\sigma_{\langle \Delta r_{l,l+2} \rangle}$ (%)	Z	$\sigma(Z)$	$\sigma(Z)/Z$ (%)	χ^2_{N-2}
GS-Lin r_{02}	10	0.057	0.01798	± 0.00013	0.7	1.7
GS-Lin r_{13}	10	0.034	0.01774	± 0.00008	0.5	0.9
GS-Lin comb.			0.01785	± 0.00007	0.4	
AGS-Lin r_{02}	10	0.056	0.01617	± 0.00011	0.7	1.7
AGS-Lin r_{13}	10	0.050	0.01604	± 0.00011	0.7	1.9
AGS-Lin comb.			0.01611	± 0.00008	0.5	

Table 6. Effective standard deviations due to uncertainties in the radiative opacities and equation of state

Quantity	Δr_{02} (%)	Δr_{13} (%)	Z (%)	μ_c (%)
σ_{opac}	0.16	0.064	0.083	0.077
σ_{EOS}	0.53	0.28	0.047	0.031

Table 7. μ_c results for ‘Monte Carlo’ models

Set	N	$\sigma_{\langle \Delta r_{l,l+2} \rangle}$ (%)	μ_c	$\sigma(\mu_c)$	$\sigma(\mu_c)/\mu_c$ (%)
GS-Cons r_{02}	5000	0.82	0.7234	± 0.0051	0.7
GS-Cons r_{13}	5000	0.56	0.7229	± 0.0033	0.6
GS-Cons comb.			0.7231	± 0.0030	0.4
GS-Opt r_{02}	1000	0.65	0.7233	± 0.0049	0.7
GS-Opt r_{13}	1000	0.38	0.7228	± 0.0030	0.4
GS-Opt comb.			0.7230	± 0.0026	0.4
AGS-Cons r_{02}	1000	0.76	0.7214	± 0.0046	0.6
AGS-Cons r_{13}	1000	0.52	0.7206	± 0.0034	0.5
AGS-Cons comb.			0.7209	± 0.0027	0.37
AGS-Opt r_{02}	5000	0.63	0.7239	± 0.0047	0.6
AGS-Opt r_{13}	5000	0.37	0.7225	± 0.0029	0.4
AGS-Opt comb.			0.7229	± 0.0025	0.3

Table 8. Z results for ‘Monte Carlo’ models

Set	N	$\sigma_{\langle \Delta r_{l,l+2} \rangle}$ (%)	Z	$\sigma(Z)$	$\sigma(Z)/Z$ (%)
GS-Cons r_{02}	5000	1.03	0.0189	$^{+0.0059}_{-0.0045}$	28
GS-Cons r_{13}	5000	0.83	0.0186	$^{+0.0052}_{-0.0040}$	25
GS-Cons comb.			0.0187	$^{+0.0039}_{-0.0030}$	19
GS-Opt r_{02}	1000	0.72	0.0197	$^{+0.0057}_{-0.0045}$	26
GS-Opt r_{13}	1000	0.52	0.0192	$^{+0.0043}_{-0.0035}$	21
GS-Opt comb.			0.0194	$^{+0.0035}_{-0.0028}$	16
AGS-Cons r_{02}	1000	0.90	0.0233	$^{+0.0072}_{-0.0055}$	27
AGS-Cons r_{13}	1000	0.75	0.0226	$^{+0.0062}_{-0.0049}$	25
AGS-Cons comb.			0.0229	$^{+0.0047}_{-0.0037}$	18
AGS-Opt r_{02}	5000	0.72	0.0220	$^{+0.0047}_{-0.0039}$	20
AGS-Opt r_{13}	5000	0.50	0.0219	$^{+0.0038}_{-0.0032}$	16
AGS-Opt comb.			0.0219	$^{+0.0029}_{-0.0025}$	12




Fig. 1.— Upper two panels: weighted mean differences, $\langle \Delta r_{02} \rangle$ (upper left-hand panel) and $\langle \Delta r_{13} \rangle$ (upper right-hand panel), versus $\ln \mu_c$, plotted for a sequence of 10 solar models. All models have the same relative mixture of heavy elements, corresponding the Grevesse & Sauval (1998) mixture but differ in the total amount of metals. The solid line in each panel is the best-fitting straight line. The dotted lines intersect at the location along each best-fitting line where the weighted mean difference is zero. Lower panels: as per upper panels, but for a sequence of 10 solar models having a relative mixture of heavy elements corresponding to the Asplund et al. (2005) mixture.




Fig. 2.— Upper two panels: weighted mean differences, $\langle \Delta r_{02} \rangle$ (upper left-hand panel) and $\langle \Delta r_{13} \rangle$ (upper right-hand panel), versus $\ln Z$, plotted for a sequence of 10 solar models. All models have the same relative mixture of heavy elements, corresponding the Grevesse & Sauval (1998) mixture but differ in the total amount of metals. The solid line in each panel is the best-fitting straight line. The dotted lines intersect at the location along each best-fitting line where the weighted mean difference is zero. Lower panels: as per upper panels, but for a sequence of 10 solar models having a relative mixture of heavy elements corresponding to the Asplund et al. (2005) mixture.




Fig. 3.— The relation between Z and μ_c (the average mean molecular weight in the inner 20% by radius) for models constructed with the GS98 relative heavy element abundances and the AGS05 relative heavy element abundances.




Fig. 4.— Upper two panels: weighted mean differences, $\langle \Delta r_{02} \rangle$ (upper left-hand panel) and $\langle \Delta r_{13} \rangle$ (upper right-hand panel), versus $\ln \mu_c$, plotted for the 5000 Monte Carlo ‘GS-Cons’ solar models. These models were made with the ‘conservative’ (large) abundance uncertainties, centered on the GS98 mixture. The solid line in each panel is the best-fitting straight line. The dotted lines intersect at the location along each best-fitting line where the weighted mean difference is zero. Lower panels: As per the upper panels, but for the 1000 Monte Carlo ‘GS-Opt’ solar models. These models were made with the ‘optimistic’ (small) abundance uncertainties, centered on the GS98 mixture.




Fig. 5.— Upper two panels: weighted mean differences, $\langle \Delta r_{02} \rangle$ (upper left-hand panel) and $\langle \Delta r_{13} \rangle$ (upper right-hand panel), versus $\ln \mu_c$, plotted for the 1000 Monte Carlo ‘AGS-Cons’ solar models. These models were made with the ‘conservative’ (large) abundance uncertainties, centered on the AGS05 mixture. The solid line in each panel is the best-fitting straight line. The dotted lines intersect at the location along each best-fitting line where the weighted mean difference is zero. Lower panels: As per the upper panels, but for the 5000 Monte Carlo ‘AGS-Opt’ solar models. These models were made with the ‘optimistic’ (small) abundance uncertainties, centered on the AGS05 mixture. Although visually the solid lines on the lower plots do not look like the best-fitting lines, they are the unbiased lines given by a least-squares fit.




Fig. 6.— Upper two panels: weighted mean differences, $\langle \Delta r_{02} \rangle$ (upper left-hand panel) and $\langle \Delta r_{13} \rangle$ (upper right-hand panel), versus $\ln Z$, plotted for the 5000 Monte Carlo ‘GS-Cons’ solar models. These models were made with the ‘conservative’ (large) abundance uncertainties, centered on the GS98 mixture. The solid line in each panel is the best-fitting straight line. The dotted lines intersect at the location along each best-fitting line where the weighted mean difference is zero. Lower panels: As per the upper panels, but for the 1000 Monte Carlo ‘GS-Opt’ solar models. These models were made with the ‘optimistic’ (small) abundance uncertainties, centered on the GS98 mixture.




Fig. 7.— Upper two panels: weighted mean differences, $\langle \Delta r_{02} \rangle$ (upper left-hand panel) and $\langle \Delta r_{13} \rangle$ (lower right-hand panel), versus $\ln Z$, plotted for the 1000 Monte Carlo ‘AGS-Cons’ solar models. These models were made with the ‘conservative’ (large) abundance uncertainties, centered on the AGS05 mixture. The solid line in each panel is the best-fitting straight line. The dotted lines intersect at the location along each best-fitting line where the weighted mean difference is zero. Lower panels: As per the upper panels, but for the 5000 Monte Carlo ‘AGS-Opt’ solar models. These models were made with the ‘optimistic’ (small) abundance uncertainties, centered on the AGS05 mixture. Although visually the solid lines on the lower plots do not look like the best-fitting lines, they are the unbiased lines given by a least-squares fit.